

EDITORIAL REVIEW

Proteolytic enzymes as mediators of glomerular injury

Despite significant advances in our understanding of the pathogenesis of glomerular injury, the specific mechanisms responsible for the pathological changes in glomerular structure and function which occur in glomerular disease remain poorly understood. Several diverse lines of evidence, including studies with purified proteinases, studies with freshly isolated glomeruli and intact cells, and cells in culture, as well as recent *in vivo* studies indicate that proteolytic enzymes play important roles as mediators of glomerular injury. The concept of proteolytic enzymes as mediators of glomerular injury is not new. The ability of polymorphonuclear leukocyte (PMN)-derived proteinases to degrade glomerular basement membrane (GBM) *in vitro* coupled with the PMN-dependence of certain types of glomerulonephritis led Cochrane and coworkers in the mid-sixties to postulate that neutrophil-derived proteinases cause glomerular injury by proteolysis of the GBM [1–3]. Since then, a large number of studies have provided support for this hypothesis. In addition, recent studies documenting the presence of GBM-degrading proteinases in glomeruli, plasma, and a variety of other circulating cells indicate that GBM damage can also be mediated by proteinases originating from these sources as well. Thus proteinase-mediated GBM damage may be an important mechanism of glomerular injury in leukocyte-independent as well as leukocyte-dependent glomerular diseases. Furthermore, recent studies suggest that proteolytic enzymes may contribute to glomerular injury by mechanisms independent of GBM damage. Such mechanisms include: proteolysis of non-GBM proteins; proteinase-mediated alterations in the production of key glomerular metabolites; and proteinase-induced effects on cellular proliferation. In the present review, we attempt to bring together a large body of diverse information derived from *in vitro* and *in vivo* studies, which supports a pathogenic role for proteolytic enzymes in glomerular injury.

Proteolytic enzymes: Nomenclature and classification

Proteolytic enzymes, (peptidases) are classified as exo- or endopeptidases. Exopeptidases are proteinases which cleave peptide bonds close to the end of a polypeptide chain. Endopeptidases (also referred to as proteinases and proteases) cleave internal peptide bonds. Although many exopeptidases are present in the kidney and may play important roles in glomerular injury, most studies concerning the role of proteolytic enzymes in glomerular injury have focused on endopeptidases. In contrast to most enzymes, which are named and classified according to their substrate and the type of reaction catalyzed (such as alcohol dehydrogenase), endopeptidases are classified

according to their catalytic mechanisms and named for the essential catalytic component (usually an amino acid) in their active site [4]. Table 1 lists the four classes of endopeptidases and some general properties of each class.

Potential sources of proteolytic enzymes for glomerular injury

With few exceptions, a proteolytic enzyme must have access to the glomerulus in order to produce glomerular injury. As mentioned above, the PMN has long been recognized as a source of proteolytic enzymes available for glomerular injury [1–3, 5]. However, other circulating blood cells such as monocytes, macrophages, and platelets contain a variety of proteolytic enzymes [6] and thus must also be included as potential sources of proteolytic enzymes. In addition, recent studies from our laboratory have demonstrated that glomeruli contain several proteolytic enzymes, most notably metalloproteinases [7, 8] and cysteine proteinases [9], which may also contribute to glomerular injury. Thus the glomerulus itself, including mesangial, endothelial, and epithelial cells constitutes a second important source of proteolytic enzymes available for glomerular injury. A third potential source of proteolytic enzymes is the plasma. Well-documented proteolytic enzymes in plasma include components of the coagulation, complement, and fibrinolytic systems. Although these plasma proteinases circulate as zymogens (inactive forms) and are usually activated only as part of well-regulated physiological processes, several lines of evidence suggest that they participate in processes which may contribute to glomerular injury. For example, thrombin appears to play an important role in the intraglomerular fibrin deposition which occurs in certain types of glomerulonephritis [10]. In addition, recent studies suggest that plasmin, generated from plasma plasminogen by glomerular plasminogen activators, may also play an important role(s) in glomerular injury. Table 2 includes a list of several proteinases of potential interest to glomerular injury as well as potential sources of each of these proteinases.

It is important to recognize that interactions between proteolytic enzymes, including those from different sources, can and do occur. For example, Gavrilovic et al [11–13] have demonstrated that the degradation of type I collagen films by phorbol myristate acetate- or interleukin-1-stimulated cells in culture is dependent on the presence of plasminogen in the medium. However, collagen degradation was inhibited by the collagenase inhibitors TIMP (tissue inhibitor of metalloproteinases) and antibodies to collagenase. Based on these observations, these workers postulated that plasminogen activator(s), produced by the stimulated cells, converted plasminogen to plasmin which in turn activated procollagenase (also produced by the cells) which is directly responsible for the collagen degradation. Clearly the potential for such interactions (and others as discussed below) must be recognized in the design of experimental protocols as well as the interpretation of results.

Table 1. Classes and some general properties of endopeptidases^a

Property	Serine	Cysteine	Aspartic	Metallo
Old name	Serine	Thiol	Carboxyl	Metallo
Enzyme commission #	3.4.21	3.4.22	3.4.23	3.4.24
Active site component	Serine	Cysteine	Aspartic acid	Zn ²⁺
pH range ^b	7-9	3-7	2-6	5-9
Inhibitors (in vitro)	PMSF, SBTI, DIFP	E-64, Iodoacetate, organo-mercurials	Pepstatin	EDTA, o-phenanthroline
Inhibitors (in vivo)	Plasma proteinase inhibitors ^c	Cystatins, α_1 -cysteine proteinase inhibitor	Unknown	TIMP-1, TIMP-2
Location	Intra- and extra-cellular	Lysosomes ^d	Lysosomes	Intra- and extracellular
Latent forms	Yes	Yes	No	Yes
Examples	Elastase, plasmin	Cathepsins B & L	Cathepsin D	Gelatinase, type IV collagenase

^a Adopted from [4]^b Varies with substrate and assay conditions^c α_1 -Proteinase inhibitor, α_2 -antiplasmin, α_2 -macroglobulin, etc. [106]^d Except calpain, the cytosolic & membrane-bound Ca²⁺-dependent cysteine proteinase [155]

Table 2. Proteolytic enzymes potentially involved as mediators of glomerular injury

Proteinase	Class	Source	Proposed mechanism of glomerular injury ^a		
			Proteolysis of GBM	Proteolysis of nonGBM proteins	Altered glomerular metabolism
Elastase & cathepsin G	Serine	Leukocytes	5, 33-40, 57, 58, 60, 61	41, 42, 85, 93, 109, 112, 114, 115	23, 25, 26, 125, 131
Plasmin	Serine	Plasma	94-99	10-13, 102-104	
Thrombin	Serine	Plasma	95	10	20-22, 24, 27, 29, 138
Gelatinase	Metallo	Leukocytes	41, 56, 59-61	113	
Gelatinase	Metallo	Glomeruli	50-52		
Glomerular metalloproteinase(s)	Metallo	Glomeruli	7, 8		
Cathepsins B & L	Cysteine	Glomeruli	9, 67-71	102, 111	

^a Numbers refer to reference number

Potential mechanisms of proteinase-mediated glomerular injury

In vitro studies have delineated a variety of biological effects of proteolytic enzymes of potential relevance to glomerular injury. For purposes of discussion, we have grouped these effects into three general categories of potential mechanisms of proteinase-mediated glomerular injury: proteolysis of the GBM; proteolysis of non-GBM proteins; and proteinase-mediated alterations in glomerular metabolism, (including effects on cell proliferation). It is important to point out that these mechanisms are not mutually exclusive. As described below and listed in Table 2, the ability of several proteolytic enzymes (such as elastase, plasmin) to produce effects described in more than one category is well documented.

Proteolysis of the GBM

The GBM is the major filtration barrier of the glomerulus restricting the passage of plasma proteins into the urine [14-16]. The hallmark of glomerular disease is proteinuria resulting from an increase in the permeability of the GBM to plasma proteins. Thus, possibly the most obvious mechanism by which proteolytic enzymes can cause glomerular injury is degradation of the GBM. The GBM is composed primarily of type IV collagen, with lesser amounts of laminin, entactin (nidogen), and heparan sulfate proteoglycan. Fibronectin, collagen types V and VII,

osteonectin (SPARC, BM-40) and several other proteins are also present in GBM, although it is presently unclear whether these proteins are intrinsic GBM components or passively accumulated following GBM synthesis [17-19]. It should be noted that each component of the GBM, including heparan sulfate proteoglycan, is susceptible in vitro to proteolysis by one or more proteinases available to or present in normal glomeruli. As discussed in detail below, a large body of evidence suggests that proteinase-mediated GBM damage is an important mechanism of glomerular injury.

Proteolysis of non-GBM proteins

In addition to GBM degradation, proteolysis of non-GBM proteins can also occur. For example, inactivation of proteinase inhibitors could result in glomerular injury mediated by uncontrolled proteinase activity. In vitro the ability of proteolytic enzymes to inactivate proteinase inhibitors is well documented (see below) and raises the possibility that such inactivation may represent an additional mechanism by which proteinases could contribute to glomerular injury.

In contrast to the proteolysis of GBM and proteinase inhibitors, which results in loss of function, proteolysis of many proteins results in increased activity of the target protein. Pertinent examples are the proteinase-mediated activation of

zymogens, the inactive forms of many enzymes. Normally zymogen activation occurs by limited proteolysis, carried out by a specific proteinase as part of a well controlled physiological process, such as coagulation or fibrinolysis. However, "inappropriate" activation of zymogens does occur and can lead to pathological changes, as evidenced by intraglomerular fibrin deposition resulting from the localized activation of thrombin [10].

Thus proteolysis of non-GBM proteins, resulting in either activation or inactivation of the target protein, represents a second potential mechanism of proteinase-mediated glomerular injury.

Proteinase-mediated alterations in glomerular metabolism

The ability of proteolytic enzymes to influence the production of several key metabolites, autocrine mediators, and intracellular signalling agents is well documented. For example, addition of proteolytic enzymes to intact cells in vitro, has resulted in altered production and/or secretion of cyclic nucleotides [20, 21], prostaglandins [22–24], platelet activating factor [25], Ca^{2+} flux and inositol polyphosphate synthesis [26, 27], adenine nucleotides [23], and reactive oxygen metabolites [28]. Each of these metabolites is postulated to play an important role in glomerular pathophysiology. In addition, recent studies [27, 29] indicate that certain proteolytic enzymes stimulate the proliferation of cultured glomerular cells. Although only limited data is available documenting altered metabolism in intact glomeruli or glomerular cells, similarities in the pathways for the production and catabolism of these metabolites among various tissues and the well-documented ability of proteinases to alter metabolism in other tissues suggests that this may be an important mechanism of proteinase-mediated glomerular injury.

In vitro evidence for proteinase-mediated glomerular injury

Proteolysis of GBM and GBM components

GBM degradation in leukocyte-dependent glomerulonephritis. Leukocytes contain proteinases in each of the four classes [6] which have been shown to effectively degrade GBM in vitro under appropriate conditions. However, several lines of evidence suggest that the serine proteinases elastase and cathepsin G and the metalloproteinase gelatinase are the leukocyte proteinases most likely to contribute to GBM damage in vivo.

Elastase and cathepsin G are well characterized serine proteinases [reviewed in 30–32] present in neutrophils. In addition, proteinases with properties similar to elastase and cathepsin G have been reported in several tissues and cells types, including macrophages, monocytes, and platelets [30–32]. Three properties of elastase and cathepsin G are of particular interest with respect to their ability to damage the GBM: their neutral pH optimum which facilitates their action in neutral pH environments such as the plasma-GBM interface; their positive charge at plasma pH (pI s = 9 to 11) which facilitates electrostatic binding to the negatively charged GBM; and their relatively low molecular weights (27 to 35 kDa) which promotes penetration into the GBM lattice and possibly allows them to escape from higher molecular weight plasma proteinase inhibitors. Thus the physical properties of these proteinases combined with their broad substrate specificity (see below) render them ideal can-

didates for mediators of glomerular injury by damage to the GBM.

GBM degradation at neutral pH by neutrophil granule extracts (now known to contain large amounts of elastase and cathepsin G) was first reported by Janoff and Zeligs [5]. This observation was confirmed and extended in subsequent studies by Davies, Coles and Hughes [33], who demonstrated that lysates of human PMN or monocytes as well as highly-purified human PMN elastase and cathepsin G could effectively degrade GBM at neutral pH in vitro. In addition, neutrophil granule extracts, purified elastase, and/or cathepsin G have been shown to degrade specific components of the GBM including proteoglycans [34–36], fibronectin [37, 38]; and type IV collagen [39, 40].

Neutrophils contain two other major neutral proteinases: the metalloproteinases, collagenase and a 92-95 kDa gelatinase [41–45]. This gelatinase appears similar to the type IV collagenase present in several tumor cell lines [41, 46], but is a distinct gene product from the 72 kDa gelatinase present in connective tissue cells [47–49] and cultured mesangial cells [50–52]. Since collagenase is relatively specific for collagen types I, II, III, which are not present in significant amounts in GBM, a major role for this metalloproteinase in GBM degradation appears unlikely. However, Lubec and coworkers [53, 54] have reported increased collagenase activity in kidneys obtained from rats with antiGBM antibody disease [53] and immune complex glomerulonephritis [54].

Gelatinases (type IV collagenases) are named because of their ability to degrade gelatin (denatured type I collagen) in vitro [42–45, 47, 48]. In vivo, gelatinases are postulated to play important roles in the degradation of type IV collagen, the major protein component of basement membranes [41, 45, 50–52, 55]. The ability of highly-purified human PMN gelatinase to degrade in vitro collagen types IV and V as well as fragments of type I (generated by collagenase) is well documented [41–46, 49]. PMN gelatinase can also degrade intact GBM at neutral pH (pH range 6.0 to 8.6) as shown by studies from our laboratory [56]. Compared to gelatinase (human PMN), collagenase (human fibroblasts) was nearly inactive in the degradation of intact GBM [56].

Several studies have extended these observations using stimulated intact neutrophils. Bray, Hume and Robinson [57] have reported that rabbit neutrophils degrade GBM in vitro at neutral pH. The amount of degradation was increased when the GBM was pretreated with antiGBM antibodies. Soya bean trypsin inhibitor and alpha-1-proteinase inhibitor (serine proteinase inhibitors) inhibited GBM degradation suggesting the involvement of elastase and/or cathepsin G. Vissers, Winterbourn and Hunt [58] have examined GBM degradation by neutrophils using "model systems" consisting of intact human neutrophils incubated with isolated human GBM impregnated with either nonspecific IgG or albumin-antialbumin immune complexes. GBM degradation was markedly inhibited by the serine proteinase inhibitor phenylmethane sulfonylfluoride (PMSF), suggesting that elastase and/or cathepsin G account for most of the GBM degradation in this system. We have shown [59] that GBM degradation by phorbol myristate acetate-stimulated PMN is mediated by activation of a metalloproteinase (possibly gelatinase) which is activated by hypochlorous acid (or a similar oxidant) generated by the PMN myeloperoxidase- H_2O_2 -halide

system. Using their *in vitro* model of immune complex GBM injury (see above), Vissers and Winterbourn [60] estimated that neutrophil gelatinase accounted for approximately 30% of the GBM degradation with elastase/cathepsin G responsible for the remainder. In a subsequent study comparing GBM degradation by human neutrophils and monocytes, Vissers et al [61] observed that GBM degradation by monocytes was inhibited nearly equally by PMSF and *o*-phenanthroline, suggesting that both serine (elastase, cathepsin G) and metalloproteinases (gelatinase) were involved.

Taken together, these studies leave little doubt concerning the ability of leukocyte proteinases to degrade GBM *in vitro*. Coupled with the *in vivo* studies discussed below they provide strong support for proteinase-mediated GBM degradation as a potentially important mechanism of glomerular injury in those types of glomerular disease characterized by significant leukocyte infiltration.

GBM degradation in leukocyte-independent glomerulonephritis. Several types of glomerular disease, including both experimentally-induced and human glomerulonephritis, lack significant leukocyte involvement. Thus leukocytes can not serve as major sources of proteolytic enzymes for glomerular injury in these types of glomerular disease. Recent studies have documented the presence of GBM-degrading proteinases in freshly isolated glomeruli and cultured glomerular cells, raising the possibility that endogenous glomerular proteinases may be responsible for GBM damage in these types of glomerulonephritis. Although glomeruli contain proteinases from each major class, published studies to date suggest that endogenous glomerular metalloproteinases and cysteine proteinases may be the most important as mediators of GBM damage in leukocyte-independent GN.

Metalloproteinases. Lovett and coworkers have identified, characterized, and purified a gelatinase, which is present in the medium (but not the cells) obtained from cultured rat [50, 51] and human [52] mesangial cells. Similar to other matrix-degrading metalloproteinases, both the rat and human mesangial cell gelatinase are synthesized and secreted as inactive zymogens which can be activated *in vitro* by aminophenyl mercuric acetate (APMA) or limited proteolysis with trypsin [50–52]. The physiological activator(s) of latent gelatinase are unknown. The purified gelatinase from either species: had a molecular weight of 66 to 68 kDa; was optimally active at neutral pH; and degraded intact GBM, soluble type IV collagen, and gelatin but not fibronectin, albumin, proteoglycan, or interstitial collagen (types I, II, and III). These data suggest that the mesangial cell gelatinase is similar to the 72 kDa gelatinase found in connective tissue cells [47–49]. These workers also reported [52] that cultured human mesangial cells secrete a protein inhibitor of gelatinase with the characteristics of TIMP-1 (tissue inhibitor of metalloproteinases). TIMP-1 [62] and TIMP-2 [63–65] are functionally similar but distinct gene products thought to play major roles in the physiological control of gelatinase activity [11–13, 62]. Lovett and coworkers reported that immunohistochemical staining of glomeruli demonstrated the presence of both gelatinase and TIMP in the mesangial area of human biopsy specimens [52].

Studies from our laboratory [7, 8] have documented that

glomeruli isolated from normal rats also contain one or more GBM-degrading metalloproteinases. Incubation of freshly isolated rat glomeruli with intact GBM at pH 7.5 resulted in significant GBM degradation as measured by hydroxyproline release [7]. GBM degradation was markedly inhibited by the metalloproteinase inhibitors, EDTA and *o*-phenanthroline, but unaffected by inhibitors of serine (PMSF, leupeptin, α -1-Pi, SBTI) or cysteine proteinases (E-64, leupeptin), documenting that metalloproteinase activity was responsible for the GBM degradation. In a separate study [8] we have characterized a gelatin-degrading metalloproteinase activity present in freshly isolated rat glomeruli. Comparison of the properties of this metalloproteinase (membrane-associated; molecular weight = 116 to 120 kDa; lack of inhibition by TIMP, lack of activation by APMA) with those of other well characterized matrix metalloproteinases (such as, gelatinase, type IV collagenase, stromelysin) indicates that this glomerular metalloproteinase is distinct from these metalloproteinases including the mesangial cell gelatinase [50–52]. Whether the GBM and gelatin degrading proteinases present in isolated rat glomeruli are the same or separate enzymes is unclear at the present time.

Although direct evidence linking endogenous glomerular metalloproteinases to GBM damage (or other types of glomerular injury) *in vivo* has not been reported, their ability to act at neutral pH and to degrade GBM suggests an important role for these proteinases in glomerular pathophysiology.

Cysteine proteinases. Cysteine proteinases are a diverse class of lysosomal enzymes with a relatively broad substrate specificity and acid pH optima [66]. Among the best characterized of the cysteine proteinases are Cathepsins B, H, and L. The ability of highly purified cathepsins B and L to degrade intact GBM *in vitro* has been reported by our lab [67] as well as by Davies and coworkers [68–70]. In our studies [67], cathepsin L was more effective in GBM degradation than several other proteinases. In addition to intact GBM, purified cysteine proteinases have been shown *in vitro* to degrade purified GBM components including collagens types IV, and V (B and L: [69, 70], proteoglycan (B, H, and L: [69, 71]; and laminin, (B and L: [69]).

Several lines of evidence, including enzyme activity [9, 72], immunohistochemistry [73, 74], and fluorescence microscopy [9], have established the presence of cathepsins B, H, and/or L in glomeruli isolated from normal rats. Furthermore, we have recently demonstrated the ability of isolated rat glomeruli to utilize endogenous cysteine proteinases to degrade intact GBM *in vitro* [9].

These studies document the ability of endogenous glomerular cysteine proteinases to degrade GBM and GBM components *in vitro*. However, most cysteine proteinases, including those documented to be present in glomeruli, are relatively inactive at neutral pH [9, 66, 67]. Thus special conditions (for example, an acidified microenvironment) must be envisioned for these proteinases to effectively act in the neutral pH of the glomerular capillary loop. Although such conditions have not been clarified, these data coupled with the beneficial effects of cysteine proteinase inhibitors in an experimental model of glomerular disease (see below) indicate an important role for cysteine proteinases in the pathogenesis of glomerular disease.

Urinary excretion of GBM components and proteolytic enzymes in glomerulonephritis

If proteinase-mediated GBM damage is indeed a mechanism of glomerular injury *in vivo*, one might expect to find increased amounts of GBM components and proteinase activity in the urine obtained from patients and animals with glomerulonephritis. Such results have been reported in both leukocyte-dependent and leukocyte-independent types of glomerulonephritis, including experimentally-induced antiGBM antibody disease [3, 75–77] and puromycin aminonucleoside-induced glomerulonephritis [78–81]. Similar results have also been reported for patients with various types of glomerulonephritis, including diffuse proliferative glomerulonephritis [82], systemic lupus and membranous glomerulonephritis [83], and acute post-strep glomerulonephritis [84]. Taken together, these studies provide further support for proteinase-mediated GBM damage as a potential mechanism of glomerular injury.

A summary of the evidence supporting GBM degradation as an important mechanism of proteinase-mediated glomerular injury is presented in Table 2.

Proteolysis of non-GBM proteins

In addition to the protein components of the GBM, a variety of structurally and/or functionally important intra- and extracellular proteins (for example, receptors, cytoskeletal proteins, enzymes, etc.) are susceptible to proteolysis by proteolytic enzymes. It is important to point out that proteolysis of such proteins may result in either activation or inactivation of the target protein. Of particular interest with respect to proteinase-mediated glomerular injury is the inactivation of proteinase inhibitors and the activation of latent proteolytic enzymes (zymogens). Although most of the examples presented below involve proteinases and target proteins from other tissues, the availability of similar proteinases and protein targets in glomeruli raises the possibility that these interactions may occur in glomeruli as well.

Activation of latent proteinases. The ability of proteolytic enzymes to activate other enzymes, often latent proteinases (zymogens) themselves, is well documented. Although all classes of proteolytic enzymes have been documented to activate one or more proteinases, activation by serine proteinases appears to be the most common and offers the greatest potential with respect to glomerular injury. Because of their many effects potentially relevant to glomerular injury, the activation of latent gelatinase (leukocyte or mesangial cell) and the conversion of plasminogen to plasmin are of particular interest in this regard.

Both elastase [85] and cathepsin G [42] have been reported to activate latent leukocyte gelatinase *in vitro* [see however, 86]. Although the mesangial cell gelatinase is a distinct gene product from the leukocyte gelatinase [46], the ability of a variety of agents (SDS, organomercurials, trypsin) to activate either of these gelatinases indicates similarities in their activation mechanisms and raises the possibility that elastase and/or cathepsin G may also activate latent mesangial cell gelatinase. Such interactions could represent an important mechanism for the activation of gelatinase in both leukocyte-dependent and independent types of glomerulonephritis.

Several lines of evidence suggest that plasmin, produced from plasminogen by plasminogen activators, may be an impor-

tant mediator of glomerular injury. Intraglomerular fibrin deposition is thought to play an important role in the development and progression of several types of experimental as well as human glomerulonephritis [10]. Clearly the ability of the glomerulus to influence fibrin deposition via fibrinolysis may play an important role in this process. The presence of fibrinolytic activity in the glomerulus is well documented and apparently mediated by the conversion of plasminogen to plasmin [87–89]. Both tPA and uPA type plasminogen activators have been identified in glomerular endothelial, epithelial and/or mesangial cells [90–92]. In addition, Machovich and Owen [93] have reported that neutrophil elastase enhances the rate of plasminogen activation by two or more orders of magnitude by a combination of effects involving increased susceptibility of plasminogen to plasminogen activators and inactivation of alpha-2-antiplasmin.

In addition to its role in fibrinolysis, plasmin has other effects of potential relevance to glomerular injury. The ability of plasmin to degrade components of the GBM, including proteoglycan [94]; the glycoproteins laminin and fibronectin [95–98]; and type IV collagen [99] has also been reported. Jones and coworkers [100, 101] have reported that glycoproteins markedly inhibit the degradation of collagen and elastin present in the extracellular matrix secreted by smooth muscle cells in culture and that plasmin, by degrading these glycoproteins, accelerated the rate of collagen and elastin degradation. Thus plasmin may accelerate the degradation of GBM collagen (by gelatinase or elastase for example) by removal of the glycoprotein components of GBM. In addition to these effects, plasmin may activate latent gelatinase. Based on the ability of plasmin to activate latent collagenase [11–13, 102–104] and apparent similarities in the activation mechanisms of latent gelatinase and latent collagenase by other agents [105], the possibility that plasmin may activate latent gelatinase must also be considered. However, Nagase and coworkers [86] failed to observe activation of latent fibroblast gelatinase by purified plasmin.

The high concentration of plasminogen in the plasma, its continuous presence in the glomerulus, the production of plasminogen activators by glomerular cells, and the ability of plasmin to produce a variety of biological effects potentially related to glomerular injury suggest an important role for this proteinase as a mediator of glomerular injury.

Inactivation of proteinase inhibitors. Several inhibitors of proteolytic enzymes are well characterized proteins which are thought to play important roles in the regulation of proteinase activity *in vivo*. Such inhibitors include the plasma proteinase inhibitors such as alpha-1-proteinase inhibitor, alpha-2-antiplasmin, alpha-1-cysteine proteinase inhibitor, etc. [reviewed in 106], the cystatins, intracellular inhibitors of cysteine proteinases [reviewed in 107], and TIMP (tissue inhibitor of metalloproteinases) [63–65, reviewed in 62]. As proteins, these inhibitors are susceptible to degradation by proteolytic enzymes. A good example of the importance of proteinase inhibitors *in vivo* is demonstrated by the high incidence of emphysema, apparently mediated by neutrophil elastase, which occurs in patients with severe alpha-1-proteinase inhibitor (alpha-1-PI) deficiency [108]. Clearly, decreased levels of proteinase inhibitors (either systemic or localized to the pericellular microenvironment) potentially could result in glomerular injury via excess proteolytic enzyme activity. Proteolytic enzymes have been reported

to inactivate several proteinase inhibitors including C1 inactivator [109] and antithrombin III [110]. However, based on the effects of elastase, gelatinase, and plasmin on GBM components (as discussed above) and on the production of key metabolites (see below), inactivation of the inhibitors of these proteinases (alpha-1-PI, TIMP, and alpha-2-antiplasmin, respectively) is of particular interest here. Proteolytic inactivation of alpha-1-PI in vitro by cathepsin L [111], macrophage elastase [112], and a metalloproteinase secreted by PMA-stimulated human neutrophils [113] has been reported. Okada et al [114] have reported that in vitro human neutrophil elastase, trypsin, and chymotrypsin (but not cathepsin G, pancreatic elastase, or plasmin) effectively inactivate TIMP-1, a major physiological inhibitor of gelatinase and other matrix metalloproteinases [11–13, 62–65]. Neutrophil elastase has been reported to inactivate alpha-2-antiplasmin [109, 115]. That these or similar reactions take place in vivo is indicated by the observation that proteolyzed alpha-1-PI has been recovered from the sites of inflammation [116, 117]. Although the pathophysiological significance of these observations to glomerular injury remains to be established, the presence of several of these proteinases in glomeruli (either intrinsically or via infiltrating leukocytes) raises the possibility that inactivation of one or more proteinase inhibitors could result in increased activity of the corresponding proteinase with subsequent deleterious effects on glomerular structure and function. In keeping with this idea, it is interesting to note that several cases of glomerulonephritis associated with severe genetic deficiency of alpha-1-PI have been reported [118–121].

The ability of proteinases to activate latent proteinases and inactivate the corresponding proteinase inhibitors raises the possibility of a potentially damaging “synergism” as a result of these combined effects. For example, elastase has been reported to activate latent gelatinase [85] and to inactivate the major physiological gelatinase inhibitor, TIMP [41, 114]. Should these actions occur simultaneously in the same microenvironment, the net result would be a dramatic increase in the local activity of gelatinase and amplification of any effects (positive or negative) arising from the active enzyme. Clearly similar synergistic interactions can be postulated for other proteinase/proteinase inhibitor pairs such as plasmin and alpha-2-antiplasmin, and elastase and alpha-1-PI. Furthermore, it should be pointed out that such synergistic interactions can be effected by agents other than proteinases. For example, intact glomeruli, leukocytes, and cultured mesangial cells produce reactive oxygen metabolites (ROM) [reviewed in 122], agents which have been reported to activate latent gelatinase [123] and inactivate TIMP-1 [124].

A summary of the evidence supporting a role for proteinase-mediated degradation of non-GBM proteins as a mechanism of glomerular injury is presented in Table 2.

Proteinase-mediated alterations in glomerular metabolism and function

In vitro, proteolytic enzymes have been shown to alter the production by intact cells of a variety of key metabolites, autocrines, and immune mediators including: stimulation of the production of PAF (platelet activating factor) by elastase and cathepsin G [25]; stimulation of the release of adenosine nucleotides by elastase [23]; stimulation of the synthesis of PGI₂ by thrombin, trypsin, and cathepsin G [22, 23]; stimulation of PGE₂ production by elastase [125]; stimulation of Ca²⁺ flux and IP₃ (inositol trisphosphate) synthesis by elastase [26] and thrombin [27]; and inhibition of thromboxane B₂ synthesis by thrombin [24]. Thrombin has been reported to also stimulate the production of uPA, tPA, and plasminogen activator inhibitor by cultured glomerular epithelial cells [29] and to increase the expression of endothelin in cultured mesangial cells [126]. However, of particular interest here is the ability of proteinases to modulate the production of cAMP and reactive oxygen metabolites (ROM), and to stimulate cell proliferation.

Proteolytic enzymes and the cAMP system. Several recent studies have suggested an important role for cyclic nucleotides in glomerular pathophysiology [127, 128]. In addition, extensive experimental evidence indicates that cAMP/cGMP may modulate inflammatory and immune responses in a variety of tissues [129, 130]. The ability of proteolytic enzymes, including trypsin, elastase, chymotrypsin, and acrosin to alter in vitro the activity of enzymes associated with the metabolism of cAMP and cGMP is well documented [131–137]. In addition, trypsin and thrombin have been reported to increase the cAMP content of intact fibroblasts [20] and lymphocytes [136]; and thrombin increases the cGMP content of cultured neuroblastoma cells [21]. These changes in cyclic nucleotide metabolism require active proteinases, are irreversible, and do not appear to result from inactivation of phosphodiesterases. Although the precise mechanism of action is presently unclear, cleavage of an inhibitory G protein has been postulated in at least one case [135]. Similar effects have been observed in freshly isolated glomeruli. Shah [138] has reported that thrombin and trypsin caused a marked increase in the cAMP content of isolated glomeruli. Thrombin had no effect on the cGMP content of glomeruli and no effect on either cAMP or cGMP content of tubules. Thrombin did not affect the cAMP phosphodiesterase activity of the glomeruli. The glomerular response to DIFP-treated thrombin was markedly reduced suggesting that proteolytic activity of thrombin is required for the response.

Role of proteolytic enzymes in the production of reactive oxygen metabolites. Several lines of evidence suggest that reactive oxygen metabolites (ROM) may play an important pathogenic role in leukocyte-dependent as well as leukocyte-independent types of glomerulonephritis [122]. Leukocytes, including PMN, monocytes, and macrophages are known to produce ROM in response to a variety of stimuli [139]. The ability of proteinase inhibitors to inhibit ROM production by stimulated leukocytes suggests an important role for proteolytic enzymes in this process. For example, the serine proteinase inhibitors PMSF, TLCK, TPCK, aprotinin, and SBTI all markedly inhibited ROM production by human PMN and monocytes stimulated with cytochalasin E or concanavalin A [140]. In addition, pretreatment of macrophages with proteolytic enzymes (elastase, trypsin, or and cathepsin G) resulted in enhanced ROM production following stimulation of the pretreated cells [141, 142].

Isolated glomeruli [143] as well as cultured mesangial cells [144] have also been shown to generate ROM. Although a role for proteinases in glomerular ROM production has not been reported, the similarities between ROM production by glomeruli and phagocytic cells suggest that proteinases may also play

a key role in glomerular ROM production. Indeed, Basci and Shah [28] have reported that treatment of freshly isolated glomeruli with trypsin or chymotrypsin results in the production of chemiluminescence, a phenomenon usually associated with ROM. In contrast to glomeruli, tubules had only a minor chemiluminescence response. Thrombin, bacterial collagenase and plasmin had no effect on ROM production by glomeruli.

Proteinase-mediated effects on cell proliferation. Several types of human and experimental glomerular diseases are characterized by abnormal proliferation of glomerular cells. Although the mechanisms responsible for such proliferation are unclear, recently published studies document the ability of thrombin to stimulate the proliferation of cultured glomerular cells. He et al [29] have reported that human alpha-thrombin promoted the proliferation of cultured human glomerular epithelial cells. This effect was inhibited by hirudin (which blocks both receptor binding and catalytic activity of thrombin) and DIPF (an irreversible thrombin inhibitor which does not block receptor binding) indicating that both receptor binding and catalytic activity are required to produce the effect. Abboud and coworkers [27] have reported that bovine thrombin caused a dose dependent increase in DNA synthesis (^3H -thymidine uptake) in cultured human mesangial cells. This effect was blocked by hirudin and also observed with a highly purified alpha-thrombin, suggesting that contaminating proteinases were not responsible for the stimulation of proliferation. The authors did not examine the effects of DIPF-treated thrombin. Thus the requirement for receptor binding, catalytic activity or both cannot be assessed.

Although only a limited number of studies have been carried out with isolated glomeruli or glomerular cells, these studies document the ability of proteolytic enzymes to modulate the production of cAMP and ROM (in both leukocytes and glomeruli) and to stimulate the proliferation of glomerular cells in culture and suggest that this may be an important mechanism of proteinase-mediated glomerular injury. A summary of the evidence supporting a role for proteinase-mediated alterations in glomerular metabolism is presented in Table 2.

In vivo studies supporting a pathogenic role for proteolytic enzymes in experimental models of glomerulonephritis

As discussed above, in vitro studies have documented several mechanisms by which proteolytic enzymes may potentially contribute to glomerular injury. Although such studies are essential to our understanding of the pathogenesis of glomerular injury, they do not *establish* a direct pathogenic role for proteolytic enzymes in glomerular disease. In contrast, recent in vivo studies involving leukocyte-dependent and leukocyte-independent models of glomerulonephritis provide direct evidence for a pathogenic role of proteolytic enzymes in glomerular injury. However, because of the inherent complexity of in vivo systems, neither the specific proteinases nor their mechanism(s) of action have been established. Regardless of the mechanism(s), in vivo studies with experimentally-induced glomerulonephritis provide strong support for a pathogenic role for proteolytic enzymes in both neutrophil-dependent and neutrophil-independent glomerular disease.

Renal infusion of elastase or cathepsin G causes proteinuria in normal rats

One obvious corollary of the proteinase-mediated glomerular injury hypothesis is that exogenous administration of the appropriate proteolytic enzyme(s) should result in glomerular injury. Johnson et al [145] have reported that infusion of microgram quantities of either elastase or cathepsin G into the renal artery of rats resulted in marked proteinuria in the 24 hour period following the infusion. Only infusion of active elastase caused proteinuria although both active and irreversibly inactivated elastase localized equally in the glomerular capillary wall. Rats perfused with cationized IgG failed to develop proteinuria, establishing that neutralization of the glomerular anionic charge by the positively charged proteinases (pI 9 to 11) did not account for the proteinuria. Interestingly, glomeruli from kidneys perfused with active elastase were histologically normal with no foot process fusion and no evidence of inflammatory cell involvement or endothelial cell damage. Based on these data and the well-documented ability of elastase and cathepsin G to degrade GBM and GBM components in vitro (see above), the authors postulated that the proteinuria arises as a result of GBM damage by the infused proteinases. While a contribution from other effects of elastase and cathepsin G (as described above) can not be ruled out, this study documents the ability of exogenously administered proteinases to induce proteinuria in rats and provides strong support for proteinase-mediated GBM degradation as a mechanism of glomerular injury.

Beige mice do not develop antiGBM antibody-induced proteinuria

A second corollary of the proteinase-mediated glomerular injury hypothesis is that animals which lack a putative glomerular-damaging proteinase should not be susceptible to experimental models of glomerulonephritis involving that proteinase. Such evidence has recently been provided by Schrijver et al [146]. These workers compared glomerular injury and proteinuria resulting from experimentally-induced antiGBM antibody disease in control and Beige mice, the latter whose PMNs are deficient in elastase and cathepsin G. Injection of antiGBM antibodies (IgG fraction) into control mice resulted in a dose-dependent proteinuria during the 24 hour period immediately following the injection. In contrast, Beige mice injected with identical doses of antiGBM antibodies failed to develop proteinuria except at the highest dose of antiGBM IgG tested. With this dose, they excreted four times as much urinary protein as uninjected controls but still only 5% of that excreted in the antiGBM-treated normal mice. Immunofluorescence microscopy showed similar deposition of rabbit IgG and mouse complement component C3 in glomeruli from both groups of mice following treatment with antiGBM IgG. The failure of Beige mice to develop proteinuria was not due to lack of infiltrating PMN. EM demonstrated similar degrees of PMN attachment to the GBM as well as swelling and necrosis of endothelial cells in both groups. PMA-stimulated superoxide production was similar in PMN from Beige and control mice. Catalase (a scavenger of H_2O_2) and deferoxamine (an iron chelator which prevents the formation of hydroxyl radical) had no effect on the proteinuria in control mice treated with antiGBM IgG, indicating that reactive oxygen metabolites are

not involved in the development of proteinuria in this model. This study clearly demonstrates that Beige mice do not develop significant proteinuria during the heterologous phase of experimentally-induced antiGBM antibody disease despite a similar degree of PMN infiltration, and antibody and C3 deposition in their glomeruli. These latter observations are interesting since they demonstrate that deposition of antibody and complement as well as the presence of PMN is not, in itself, sufficient to cause proteinuria. These data, in combination with the *in vitro* and *in vivo* studies presented above, provide additional support for proteinase-mediated GBM damage as a mechanism of proteinuria in the neutrophil-dependent antiGBM antibody disease model of glomerulonephritis.

Effect of synthetic proteinase inhibitors on pathological changes in experimental glomerulonephritis

An additional corollary of the proteinase-mediated glomerular injury hypothesis is that inhibitors of putative glomerular-damaging proteolytic enzymes should block or attenuate the pathological changes associated with experimental models of glomerulonephritis involving those proteinases. Such studies have been reported for both leukocyte-dependent and leukocyte-independent models of glomerulonephritis.

Cysteine proteinase inhibitors reduce proteinuria in a neutrophil- and complement-independent model of antiGBM antibody disease. Studies from our laboratory have provided evidence that cysteine proteinases may play a pathogenic role in a complement- and neutrophil-independent model of antiglomerular basement membrane (GBM) antibody disease [72, 147]. Transepoxysuccinyl-L-leucylamido-(4-guanidino)butane (E-64) is a specific, irreversible inhibitor of cysteine proteinases. It is: unreactive with serine, aspartic, or metalloproteinases; does not react at appreciable rates with low molecular weight thiols such as glutathione; does not react with nonproteolytic enzymes which contain a cysteine at their active site [148, 149]; and is relatively nontoxic [150]. In two separate studies, rats treated with E-64 and proteinuric doses of neutrophil- and complement-independent antiGBM IgG excreted significantly less urinary protein than rats receiving identical doses antiGBM IgG alone (E-64-treated: -43%, $P < 0.001$). This reduction in urinary protein excretion was accompanied by a marked decrease in the specific activity of cathepsins B and L in glomeruli and cortex isolated from rats treated with antiGBM IgG and E-64 compared to either saline-treated controls or rats treated with antiGBM IgG alone. Renal function, as assessed by BUN and plasma creatinine measurements, was unchanged in either experimental group compared to saline-treated controls.

In a subsequent study [151] we confirmed and extended these observations. Rats treated with trans-epoxysuccinyl-L-leucylamido-(3-methyl)butane (Ep475, an E-64 analog) and proteinuric doses of neutrophil- and complement-independent antiGBM IgG exhibited significantly reduced proteinuria (-45%, $P < 0.05$) compared to rats receiving identical doses of antiGBM IgG only. There was a significant reduction in the activity of both cathepsin B and cathepsin L in renal cortices obtained from Ep-475-treated rats compared to either saline-treated controls or rats treated with antiGBM IgG only. Administration of Z-Phe-Tyr(O-t-butyl)CHN₂ (ZFYCHN₂), a specific, irreversible inhibitor mainly of cathepsin L [152], caused a similar

reduction in proteinuria (-56%, $P < 0.05$). This reduction in proteinuria was accompanied by a marked decrease (-84%) in the specific activity of renal cortical cathepsin L in ZFYCHN₂-treated rats. There was no significant change in: the activity of renal cortical cathepsin B; renal antiGBM antibody uptake; plasma urea nitrogen or plasma creatinine values in the ZFYCHN₂-treated rats compared to rats treated with antiGBM IgG only or saline-treated controls.

The mechanism(s) by which cysteine proteinases may contribute to glomerular injury is presently unclear. As discussed above, cysteine proteinases are present in normal glomeruli and can degrade GBM and GBM components *in vitro* under acid conditions. The extracellular location of the GBM suggests that the initial step(s) in GBM degradation must take place at the neutral pH presumed to exist at the GBM-plasma interface. Most cysteine proteinases, including cathepsins B and L, are relatively inactive *in vitro* at neutral pH, although we have observed significant degradation of GBM by endogenous glomerular cysteine proteinases at pH 6.5 [9]. However, it must be kept in mind that determination of pH optima *in vitro* are dependent on the substrate used as well as the specific conditions of the assay [153]. Furthermore, the ability of certain cells to acidify their pericellular environment [154] may provide a mechanism for lysosomal acid proteinases to function in an extracellular environment. Although calpain [155] and cathepsin S [156] are cysteine proteinases which are active at neutral pH, little is known concerning the presence of these enzymes in glomeruli or the potential role of these proteinases in GBM damage or other types of glomerular injury. Thus while a direct role for cysteine proteinases in GBM degradation remains a possibility, other mechanisms of cysteine proteinase-mediated glomerular injury must be considered including inactivation of proteinase inhibitors [111] and activation of latent proteinases as previously demonstrated for cathepsin B [102]. Regardless of the mechanism, the ability of specific cysteine proteinase inhibitors to reduce proteinuria in an experimental model of glomerulonephritis indicates an important role for one or more cysteine proteinases in this model of glomerular injury.

Serine proteinase inhibitors. Jennette et al [157] have reported a significant reduction (-69% to -89%) in glomerular necrosis in mice with horse apoferritin (HAF)-induced immune complex glomerulonephritis that had been treated with amidine-type inhibitors of trypsin-like proteinases (but which did not effectively inhibit the complement system). There was no difference between treated and non-treated groups in the amount or distribution of IgG, HAF, or C3, as measured by immunofluorescence staining. In long-term survival studies (33 days) only 3 of 9 inhibitor-treated mice died compared with 7 out of 10 mice in the group receiving HAF only.

Ikehara et al [158] have examined the effect of FUT-175 (6-amidino-2-naphthyl p-guanidinobenzoate dimethanesulphonate), a synthetic proteinase inhibitor of C1r and C1 esterase, thrombin, plasmin, kallikrein, and trypsin, on the development of lupus nephritis in (NZBxNZW)_{F1} mice. High doses of FUT-175 (400 mg/kg/day) decreased the incidence of severe proteinuria (>300 mg/dl), improved renal function (as indicated by BUN), and decreased immune complex and C3 deposits and glomerular damage as assessed histologically. FUT-175 had no effect on ANA titer. Although these inhibitors are specific for serine proteinases, the fact that they inhibit several serine

Table 3. In vivo evidence for proteinase-mediated glomerular injury

Model	Animal	End point	Proteinase	Reference
Proteinase infusion	Normal rats	Proteinuria	Elastase and cathepsin G	145
AntiGBM antibody disease ^a	Beige mice	Proteinuria	Elastase and cathepsin G	146
AntiGBM antibody disease ^b	Normal rats	Proteinuria	Cathepsin L and cathepsin B	72, 147, 151
Immune complex GN	Normal mice	Glomerular necrosis	Serine proteinases	157
Lupus nephritis	(NZBxNZW)F ₁ mice	Proteinuria, BUN, IC and C3 deposition	Serine proteinases	158

^a Neutrophil-dependent model^b Neutrophil-independent model

proteinases precludes identification of the specific serine proteinase involved or the mechanism of injury.

A summary of the in vivo evidence supporting proteinase-mediated glomerular injury is presented in Table 3.

Conclusions and perspective

A large number of in vitro studies provide strong support for the concept that proteinase-mediated GBM damage may be an important mechanism of glomerular injury in leukocyte-dependent as well as leukocyte-independent types of glomerulonephritis. In addition, recent in vitro studies indicate that proteolytic enzymes may contribute to glomerular injury by mechanisms independent of GBM damage, including activation of latent enzymes (such as, plasminogen, metalloproteinases), inactivation of proteinase inhibitors (alpha-1-proteinase inhibitor, alpha-2-antiplasmin, and tissue inhibitor of metalloproteinases), and proteinase-mediated alterations in glomerular metabolism (cAMP, reactive oxygen metabolites, Ca²⁺ and IP₃) and cell proliferation. It is important to point out that most in vitro studies have been carried out in simplified systems which lack proteinase inhibitors and other potentially important modulators known to be present in vivo. Nonetheless, in vitro studies clearly document the potential of proteolytic enzymes to mediate glomerular injury in vivo and aid in the design and interpretation of in vivo experiments.

In vivo studies utilizing experimental models of glomerulonephritis also support an important role for proteolytic enzymes in both leukocyte-dependent and independent models of glomerular disease. The potential for synergistic interactions among proteinases, proteinase inhibitors, and other effectors (for example, reactive oxygen metabolites) complicates the interpretation of in vivo studies and precludes in most cases the definitive identification of the specific proteinase involved and the mechanism(s) by which it produces glomerular injury. Nonetheless, these studies are in good agreement with the in vitro studies and provide support for the proteinase-mediated glomerular injury hypothesis. Clearly, further in vivo studies are needed to establish which types of glomerular diseases involve proteolytic enzymes and the identity of the proteinase(s) involved. The increasing availability of a variety of specific proteinase inhibitors suitable for experimental use in vivo should greatly facilitate such studies.

Glomerular disease is a multifaceted phenomenon resulting from interactions among diverse biological processes. Our

understanding of the role of proteolytic enzymes in these processes is just beginning. The recognition that proteolytic enzymes may mediate glomerular injury by mechanisms independent of GBM damage; that complex interactions can occur between proteolytic enzymes, their inhibitors, and other effector systems; and that glomerular-damaging proteolytic enzymes may originate from sources other than PMN represent important conceptual advances in our understanding of the potential role of proteinases in glomerular injury. These concepts should provide a useful framework for the design and interpretation of future studies and hopefully enhance our overall understanding of the pathogenesis of glomerular disease.

WILLIAM H. BARICOS and SUDHIR V. SHAH¹
New Orleans, Louisiana, USA

Acknowledgments

The work cited from the authors' laboratories was sponsored by grants from the American Heart Association (National Center and Louisiana Affiliate), the Research Service of the Veterans Administration Hospital and NIH grant R01-DK41480. Drs. Baricos and Shah are former Established Investigators of the American Heart Association and portions of the work cited from our laboratories was carried out during the tenure of these awards. The authors acknowledge Dr. Gillian Murphy, Strangeways Research Laboratory, Cambridge, England, for helpful discussion and critical review of the manuscript; Quoc Le, Alvin Wong, and Shirley Cortez for proofreading the manuscript; and Jill Barbay for preparation of the Tables.

Reprint requests to Dr. William H. Baricos, Department of Biochemistry, Tulane Medical School, 1430 Tulane Ave., New Orleans, Louisiana 70112, USA

References

1. COCHRANE CG, UNANUE ER, DIXON FJ: A role of polymorphonuclear leukocytes and complement in nephrotoxic nephritis. *J Exp Med* 122:99-116, 1965
2. COCHRANE CG, AIKIN BS: Polymorphonuclear leukocytes in immunological reactions. *J Exp Med* 124:733-752, 1966
3. HAWKINS D, COCHRANE CG: Glomerular basement membrane damage in immunologic glomerulonephritis. *Immunol* 14:665-681, 1968
4. BARRETT AJ: Introduction to the history and classification of

¹ Present address: Dept of Medicine, University of Arkansas for Medical Sciences, Little Rock, Arkansas 72205, USA.

- tissue proteinases, in *Proteinases in Mammalian Cells and Tissues*, edited by BARRETT AJ, Amsterdam, North-Holland, 1977, p. 1
5. JANOFF A, ZELIGS JD: Vascular injury and lysis of basement membrane in vitro by neutral protease of human leukocytes. *Science* 161:702-704, 1968
 6. HENSON PM, HENSON JE, FITTSCHEN C, KIMANI G, BRATTON DL, RICHES DWH: Phagocytic cells: degranulation and secretion, chapter 22 in *Inflammation: Basic Principles and Clinical Correlates*, edited by GALLIN JJ, GOLDSTEIN IM, SNYDERMAN R, New York, Raven Press, Ltd., 1988, p. 363
 7. NGUYEN HH, BARICOS WH, SHAH SV: Degradation of glomerular basement membrane by a neutral metalloproteinase(s) present in glomeruli isolated from normal rat kidney. *Biochem Biophys Res Commun* 141:898-903, 1986
 8. LE QC, SHAH SV, NGUYEN HH, CORTEZ SL, BARICOS WH: A novel metalloproteinase present in freshly isolated rat glomeruli. *Am J Physiol (Renal)* 260:F555-F561, 1991
 9. BARICOS WH, CORTEZ SL, LE QC, ZHOU Y, DICARLO RM, O'CONNOR SE, SHAH SV: Glomerular basement membrane degradation by endogenous cysteine proteinases present in isolated rat glomeruli. *Kidney Int* 38:395-401, 1990
 10. HANCOCK W, ATKINS R: Activation of coagulation pathways and fibrin deposition in human glomerulonephritis. *Sem Nephrol* 5:69-77, 1985
 11. GAVRILOVIC J, HEMBRY RM, REYNOLDS JJ, MURPHY G: Tissue inhibitor of metalloproteinases (TIMP) regulates extracellular type I collagen degradation by chondrocytes and endothelial cells. *J Cell Sci* 87:357-362, 1987
 12. GAVRILOVIC J, MURPHY G: The role of plasminogen in cell-mediated collagen degradation. *Cell Biol Intl Repts* 13:367-375, 1989
 13. GAVRILOVIC J, REYNOLDS JJ, MURPHY G: Inhibition of type I collagen film degradation by tumor cells using a specific antibody to collagenase and the specific tissue inhibitor of metalloproteinases (TIMP). *Cell Biol Intl Repts* 9:1097-1107, 1985
 14. VENKATACHALAM MA, RENNKE HG: The structural and molecular basis of glomerular filtration. *Circ Res* 43:377-347, 1978
 15. BRENNER BM, HOSTETTER TH, HUMES DA: Glomerular permselectivity: barrier function based on discrimination of molecular size and charge. *Am J Physiol* 234:F455-F460, 1978
 16. FARQUHAR MG: The glomerular basement membrane: a selective macromolecular filter, in *Cell Biology of the Extracellular Matrix*, edited by HAY ED, New York, Plenum, 1981, p. 335
 17. TIMPL R: Structure and biological activity of basement membrane proteins. *Eur J Biochem* 180:487-502, 1989
 18. YURCHENCO PD, SCHITTNY JC: Molecular architecture of basement membranes. *FASEB J* 4:1577-1590, 1990
 19. MARTINEZ-HERNANDEZ A, AMENTA PS: The basement membrane in pathology. *Lab Invest* 48:656-677, 1983
 20. GORDON EA, FENTON JW, CARNEY DH: Thrombin-receptor occupancy initiates a transient increase in cAMP levels in mitogenically responsive hamster (NIL) fibroblasts. *Ann NY Acad Sci* 485:249-263, 1986
 21. SNIDER RM, RICHELSON E: Thrombin stimulation of guanosine 3', 5'-monophosphate formation in murine neuroblastoma cells (clone N1E-115). *Science* 221:566-568, 1983
 22. WESKLER BB, LEY CW, JAFFE EA: Stimulation of endothelial cell prostacyclin production by thrombin, trypsin, and the ionophore A 23187. *J Clin Invest* 62:923-930, 1978
 23. LEROY EC, AGER A, GORDON JL: Effects of neutrophil elastase and other proteinases on porcine aortic endothelial prostaglandin I₂ production, adenine nucleotide release, and responses to vasoactive agents. *J Clin Invest* 74:1003-1010, 1984
 24. PODJARNY E, RATHAUS M, POMERANZ A, SHAPIRA J, BERNHEIM J: Thrombin inhibits the synthesis of prostanoids by isolated glomeruli and peritoneal macrophages in rats. *Nephron* 53:50-53, 1989
 25. CAMUSSI G, TETTA C, BUSSOLINO F, BAGLIONI C: Synthesis and release of platelet-activating factor is inhibited by plasma alpha-1-proteinase inhibitor or alpha-1-antichymotrypsin and is stimulated by proteinases. *J Exp Med* 168:1293-1306, 1988
 26. PETERSON MW, GRUENHAUPT D, SHASBY DM: Neutrophil cathepsin G increases calcium flux and inositol polyphosphate production in cultured endothelial cells. *J Immunol* 143:609-616, 1989
 27. SHULTZ PJ, KNAUSS TC, MENE P, ABBODD HE: Mitogenic signals for thrombin in mesangial cells: regulation of phospholipase C and PDGF genes. *Am J Physiol (Renal)* 257:F366-F374, 1989
 28. BASCI A, SHAH SV: Trypsin- and chymotrypsin-induced chemiluminescence by isolated rat glomeruli. *Am J Physiol* 252:C611-C617, 1987
 29. HE C-J, RONDEAU E, MEDCALF RL, LACAVE R, SCHLEUNING W-D, SRAER J-D: Thrombin increases proliferation and decreases fibrinolytic activity of kidney glomerular epithelial cells. *J Cellular Physiol* 146:131-140, 1991
 30. STARKEY PM: Elastase and cathepsin G, the serine proteinases of human neutrophil leukocytes and spleen, Chapter 1 in, *Proteinases in Mammalian Cells and Tissues*, edited by BARRETT AJ, Amsterdam, North-Holland, 1977, p. 57
 31. JANOFF A: Elastase in tissue injury. *Am Rev Med* 36:207-216, 1985
 32. HAVEMANN K, GRAMSE M: Physiology and pathology of neutral proteinases of human granulocytes. *Adv Exp Med* 167:1-20, 1984
 33. DAVIES M, COLES GA, HUGHES KT: Glomerular basement membrane injury by neutrophil and monocyte neutral proteinases. *Renal Physiol (Basel)* 3:106-111, 1980
 34. STARKEY PM, BARRETT AJ: Human cathepsin G. Catalytic and immunological properties. *Biochem J* 155:273-278, 1976
 35. KEISER H, GREENWALD RA, FEINSTEIN G, JANOFF A: Degradation of cartilage proteoglycan by human leukocyte granule neutral proteases—a model of joint injury. *J Clin Invest* 57:625-632, 1976
 36. JANOFF A, FEINSTEIN G, MALEMUD CJ, ELIAS JM: Degradation of cartilage proteoglycan by human leukocyte granule neutral proteases—a model of joint injury. *J Clin Invest* 57:615-624, 1976
 37. McDONALD JA, KELLEY DG: Degradation of fibronectin by human leukocyte elastase. *J Biol Chem* 255:8848-8858, 1980
 38. SCHMIDT W, HAVEMANN K: Isolation of elastase-like and chymotrypsin-like neutral proteinases from human granulocytes. Hoppe-Seyler's *Z. Physiol Chem* 355:1077-1082, 1974
 39. MAINARDI CL, DIXIT S, KANG AH: Degradation of type IV (basement membrane) collagen by a proteinase isolated from human polymorphonuclear leukocyte granules. *J Biol Chem* 255:5435-5441, 1980
 40. PIPOLY DJ, CROUCH EC: Degradation of native type IV procollagen by human neutrophil elastase. Implications for leukocyte-mediated degradation of basement membranes. *Biochem* 26:5748-5754, 1987
 41. WEISS SJ, PEPPIN GJ: Collagenolytic metalloenzymes of the human neutrophil: Characteristics, regulation and potential function in vivo. *Biochem Pharm* 35:3189-3197, 1986
 42. MURPHY G, BRETZ U, BAGGIOLINI M, REYNOLDS JJ: The latent collagenase and gelatinase of human polymorphonuclear neutrophil leukocytes. *Biochem J* 192:517-525, 1980
 43. MURPHY G, REYNOLDS JJ, BRETZ U, BAGGIOLINI M: Partial purification of collagenase and gelatinase from human polymorphonuclear leukocytes. *Biochem J* 203:209-221, 1982
 44. HIBBS MS, HASTY KA, SEYER JM, KANG AH, MAINARDI CL: Biochemical and immunological characterization of the secreted forms of human neutrophil gelatinase. *J Biol Chem* 260:2493-2500, 1985
 45. MURPHY G, WARD R, HEMBRY RM, REYNOLDS JJ, KUHN K, TRYGGVASON K: Characterization of gelatinase from pig polymorphonuclear leukocytes. A metalloproteinase resembling tumor type IV collagenase. *Biochem J* 258:463-472, 1989
 46. LIOTTA LA, TRYGGVASON K, GARBISA S, ROBEY PG, ABE S: Partial purification and characterization of a neutral protease which cleaves type IV collagen. *Biochem* 20:100-104, 1981
 47. SEITZER JL, ADAMS SA, GRANT GA, EISEN AZ: Purification and properties of a gelatin-specific neutral proteinase from human skin. *J Biol Chem* 256:4662-4668, 1981
 48. SAPOLSKY AI, SHEFF MF, MATSUTA K, HOWELL DS, MOSKOWITZ RW, GOLDBERG VM, NORBY DP, MALEMUD CJ: Gelatinase-like activity from articular chondrocytes in monolayer culture. *Biochim Biophys Acta* 762:227-231, 1983

49. MURPHY G, McALPINE CG, POLL CT, REYNOLDS JJ: Purification and characterization of a bone metalloproteinase that degrades types IV and V collagen. *Biochim Biophys Acta* 831:49-58, 1985
50. LOVETT DH, STERZEL RB, KASHGARIAN M, RYAN JL: Neutral proteinase activity produced in vitro by cells of the glomerular mesangium. *Kidney Int* 23:342-349, 1983
51. DAVIES M, THOMAS GJ, MARTIN J, LOVETT DH: The purification and characterization of a glomerular-basement-membrane-degrading neutral proteinase from rat mesangial cells. *Biochem J* 251:419-425, 1988
52. MARTIN J, DAVIES M, THOMAS G, LOVETT DH: Human mesangial cells secrete a GBM-degrading neutral proteinase and a specific inhibitor. *Kidney Int* 36:790-801, 1989
53. LUBEC G: Collagenase activity of rat kidney with glomerulonephritis during the heterologous phase. *Clinica Chimica Acta* 76:89-94, 1977
54. LUBEC G, RATZENHOFFER E: Collagenase activity of rat kidney with immune complex glomerulonephritis. *Clinica Chimica Acta* 82:205-207, 1978
55. MIGNATTI P, TSUBOI R, ROBBINS E, RIFKIN DB: In vitro angiogenesis on the human amniotic membrane: requirement for basic fibroblast growth factor-induced proteinases. *J Cell Biol* 108:671-682, 1989
56. BARICOS WH, MURPHY G, ZHOU Y, NGUYEN HH, SHAH SV: Degradation of glomerular basement membrane by purified mammalian metalloproteinases. *Biochem J* 254:609-612, 1988
57. BRAY J, HUME DA, ROBINSON GB: Degradation of rat or sheep glomerular basement membrane by rabbit neutrophils in vitro. *Mol Biol Med* 1:253-269, 1983
58. VISSERS MCM, WINTERBOURN CC, HUNT JS: Degradation of glomerular basement membrane by human neutrophils in vitro. *Biochim Biophys Acta* 804:154-160, 1984
59. SHAH SV, BARICOS WH, BASCI A: Degradation of human glomerular basement membrane by stimulated neutrophils. *J Clin Invest* 79:25-31, 1987
60. VISSERS MCM, WINTERBOURN CC: Gelatinase contributes to the degradation of glomerular basement membrane collagen by human neutrophils. *Collagen Rel Res* 8:113-122, 1988
61. VISSERS MCM, WIGGINS R, FANTONE JC: Comparative ability of human monocytes and neutrophils to degrade glomerular basement membrane in vitro. *Lab Invest* 60:831-838, 1989
62. CAWSTON TE: Protein inhibitors of metalloproteinases, Chapter 20 in *Proteinase Inhibitors*, edited by BARRETT AJ, SALVESEN G, Amsterdam, Elsevier, 1986, p. 589
63. GOLDBERG GI, MARMER BL, GRANT GA, EISEN AZ, WILHELM S, HE C: Human 72-kilodalton type IV collagenase forms a complex with a tissue inhibitor of metalloproteinases designated TIMP-2. *Proc Natl Acad Sci USA* 86:8207-8211, 1989
64. DE CLERCK YA, YEAN TD, RATZKIN BJ, LU HS, LANGLEY KE: Purification and characterization of two related but distinct metalloproteinase inhibitors secreted by bovine aortic endothelial cells. *J Biol Chem* 264:17445-17453, 1989
65. STETLER-STEVENSON WG, BROWN PD, ONISTO M, LEVY AT, LIOTTA LA: Tissue inhibitor of metalloproteinases-2 (TIMP-2) mRNA expression in tumor cell lines and human tumor tissues. *J Biol Chem* 265:13933-13938, 1990
66. KIRSCHKE H, LANGNER J, RIEMANN S, WIENDERANDERS B, ANSORGE S, BOHLEY P: Lysosomal cysteine proteinases, in *Protein Degradation in Health and Disease*, Amsterdam, Excerpta Medica, 1980, p. 15
67. BARICOS WH, ZHOU Y, MASON RW, BARRETT AJ: Human kidney cathepsins B and L. Characterization and potential role in degradation of glomerular basement membrane. *Biochem J* 252:301-304, 1988
68. DAVIES M, BARRETT AJ, TRAVIS J, SANDERS E, COLES GA: The degradation of human glomerular basement membrane with purified lysosomal proteinases: Evidence for the pathogenic role of the polymorphonuclear leucocyte in glomerulonephritis. *Clin Sci Mol Med* 54:233-240, 1978
69. THOMAS GJ, DAVIES M: The potential role of human kidney cortex cysteine proteinases in glomerular basement membrane degradation. *Biochem Biophys Acta* 990:246-253, 1989
70. DAVIES M, HUGHES KT, THOMAS GJ: Evidence that kidney lysosomal proteinases degrade the collagen of glomerular basement membrane. *Renal Physiol* (Basel) 3:116-119, 1980
71. MORRISON RIG, BARRETT AJ, DINGLE JT, PRIOR D: Cathepsins B1 and D. Action on human cartilage proteoglycans. *Biochim Biophys Acta* 302:411-419, 1973
72. BARICOS WH, O'CONNOR SE, CORTEZ SL, WU L-T, SHAH SV: The cysteine proteinase inhibitor, E-64, reduces proteinuria in an experimental model of glomerulonephritis. *Biochem Biophys Res Comm* 155:1318-1323, 1988
73. YOKOTA S, TSUJI H, KATO K: Immunocytochemical localization of cathepsin B in rat kidney. I. Light microscopic study using the indirect immunoenzyme technique. *J Histochem Cytochem* 34:891-897, 1986
74. YOKOTA S, TSUJI H, KATO K: Immunocytochemical localization of cathepsin B in rat kidney. II. Electron microscopic study using the protein A-gold technique. *J Histochem Cytochem* 34:899-907, 1986
75. GANG NF, TRACHTENBERG E, ALLERHAND J, KALANT N, MAUTNER W: Correlation of proteinuria, excretion of the glomerular basement membrane-like protein, and changes in the ultrastructure of the glomerular basement membrane as visualized with lanthanum. *Lab Invest* 23:436-441, 1970
76. DAVIN J-C, DAVIES M, FOIDART J-M, FOIDART JB, DECHENNE CA, MAHIEU PR: Urinary excretion of neutral proteinases in nephrotic rats with a glomerular disease. *Kidney Int* 31:32-40, 1987
77. LUBEC G, FORSTER O, CORADELLO H, MAXA E, POLLAK A: Proteolytic degradation of the glomerular basement membrane and immunochemical characterization of the split products. *Renal Physiol* 3:126-132, 1980
78. GANG NF, MAUTNER W: Studies on the mechanism of the onset of proteinuria in aminonucleoside nephrosis. *Lab Invest* 27:310-316, 1972
79. BARICOS WH, SHAH SV: Increased cathepsin D-like activity in cortex, tubules, and glomeruli isolated from rats with experimental nephrotic syndrome. *Biochem J* 223:393-399, 1984
80. MAYER M, YEDGAR S, JOFFE M, SHAFIRE E: Urine protease and antiprotease activity in experimental aminonucleoside nephrotoxicity. *Nephron* 29:223-228, 1981
81. JUUKOLA A, RISTELI J, AUTIO-HARMAINEN H, RISTELI L: Effects of experimental nephrosis on basement-membrane components and enzymes of collagen biosynthesis in rat kidney. *Biochem J* 226:243-250, 1985
82. SANDERS E, COLES GA, DAVIES M: Lysosomal enzymes in human urine: Evidence for polymorphonuclear leucocyte proteinase involvement in the pathogenesis of human glomerulonephritis. *Clin Sci Mol Med* 54:667-672, 1978
83. SHIBATA S, NAGASAWA T: Glycoproteinuria as an indicator of glomerular basement membrane damage. *J Lab Clin Med* 89:207-216, 1977
84. KARAN A, SAATCI Ü, BAKKALOGLU A: The role of cathepsin D in pathogenesis of acute poststreptococcal glomerulonephritis. *Acta Padiatr Scand* 65:355-360, 1976
85. VISSERS MCM, WINTERBOURN CC: Activation of human neutrophil gelatinase by endogenous serine proteinases. *Biochem J* 249:327-331, 1988
86. OKADA Y, MORODOMI T, ENGHILD JJ, SUZUKI K, YASUI A, NAKANISHI I, SALVESEN G, NAGASE H: Matrix metalloproteinase 2 from human rheumatoid fibroblasts. Purification and activation of the precursor and enzymic properties. *Eur J Biochem* 194:721-730, 1990
87. MYHRE-JENSEN O: Localization of fibrinolytic activity in the kidney and urinary tract of rats and rabbits. *Lab Invest* 25:403-411, 1971
88. BERGSTEIN JM, MICHAEL AF: Cortical fibrinolytic activity in normal and diseased human kidneys. *J Lab Clin Med* 79:710-719, 1972
89. SRAER JD, BOELAERT J, MIMOUNE O, MOREL-MAROGER L, HORNYCH H: Quantitative assessment of fibrinolysis on isolated glomeruli. *Kidney Int* 4:350-352, 1973
90. ANGLES-CANO E, RONDEAU E, DELARUE F, HAGEGE J, SULTAN Y, SRAER JD: Identification and cellular localization of plasmino-

- gen activators from human glomeruli. *Throm Haemos* 54:688-692, 1985
91. RONDEAU E, OCHI S, LACAVE R, HE C-J, MEDCALF R, DELARUE F, SRAER JD: Urokinase synthesis and binding by glomerular epithelial cells in culture. *Kidney Int* 36:593-600, 1989
 92. GLASS WF II, RADNIK RA, GARONI JA, KREISBERG JI: Urokinase-dependent adhesion loss and shape change after cyclic adenosine monophosphate elevation in cultured rat mesangial cells. *J Clin Invest* 82:1992-2000, 1988
 93. MACHOVICH R, OWEN WG: An elastase-dependent pathway of plasminogen activation. *Biochemistry* 28:4517-4522, 1989
 94. MOCHAN E, KELER T: Plasmin degradation of cartilage proteoglycan. *Biochim Biophys Acta* 800:312-315, 1984
 95. LIOTTA LA, GOLDFARB RH, TERRANOVA VP: Cleavage of laminin by thrombin and plasmin: Alpha thrombin selectively cleaves the beta chain of laminin. *Throm Res* 21:663-673, 1981
 96. LIOTTA LA, GOLDFARB RH, BRUNDAGE R, SIEGAL GP, TERRANOVA V, GARBISA S: Effect of plasminogen activator (urokinase), plasmin, and thrombin on glycoprotein and collagenous components of basement membrane. *Cancer Res* 41:4629-4636, 1981
 97. CARLSEN SA, RAMSHAW IA, WARRINGTON RC: Involvement of plasminogen activator with tumor metastasis in a rat model. *Cancer Res* 44:3012-3016, 1984
 98. JILEK F, HORMANN H: Cold-insoluble globulin, III. Cyanogen bromide and plasminolysis fragments containing a label introduced by transamination. *Hoppe-Seyler's Z Physiol Chem* 358:1165-1168, 1977
 99. MACKAY AR, CORBITT RH, HARTZLER JL, THORGEIRSSON UP: Basement membrane type IV collagen degradation: Evidence for the involvement of a proteolytic cascade independent of metalloproteinases. *Cancer Res* 50:5997-6001, 1990
 100. JONES PA, WERB Z: Degradation of connective tissue matrices by macrophages. II. Influence of matrix composition on proteolysis of glycoproteins, elastin, and collagen by macrophages in culture. *J Exp Med* 152:1527-1536, 1980
 101. JONES PA, DECLERCK YA: Destruction of extracellular matrices containing glycoproteins, elastin, and collagen by metastatic human tumor cells. *Cancer Res* 40:3222-3227, 1980
 102. ECKHOUT Y, VAES G: Further studies on the activation of procollagenase, the latent precursor of bone collagenase. Effects of lysosomal cathepsin B, plasmin and kallikrein, and spontaneous activation. *Biochem J* 166:21-31, 1977
 103. CAWSTON TE, MERCER E, TYLER JA: The activation of latent pig synovial collagenase. *Biochim Biophys Acta* 657:73-83, 1981
 104. WERB Z, MAINARDI CL, VATER CA, HARRIS ED: Endogenous activation of latent collagenase by rheumatoid synovial cells. Evidence for a role for plasminogen activator. *N Engl J Med* 296:1017-1023, 1977
 105. VAN WART HE, BIRKEDAL-HANSEN H: The cysteine switch: A principle of regulation of metalloproteinase activity with potential applicability to the entire matrix metalloproteinase gene family. *Proc Natl Acad Sci USA* 87:5578-5582, 1990
 106. TRAVIS J, SALVESEN GS: Human plasma proteinase inhibitors. *Ann Rev Biochem* 52:655-709, 1983
 107. BARRETT AJ, RAWLINGS ND, DAVIES ME, MACHLEIDT W, SALVESEN G, TURK V: Cysteine proteinase inhibitors of the cystatin superfamily, Chapter 18, in *Proteinase Inhibitors*, edited by BARRETT AJ, SALVESEN G, Amsterdam, Elsevier, 1986, p. 515
 108. JANOFF A: Elastases and emphysema. Current assessment of the protease-antiprotease hypothesis. *Am Rev Res Dis* 132:417-433, 1985
 109. BROWER MS, HARPEL PC: Proteolytic cleavage and inactivation of alpha-2-plasmin inhibitor and C₁ inactivator by human polymorphonuclear leukocyte elastase. *J Biol Chem* 257:9849-9854, 1982
 110. JOCHUM M, LANDER S, HEIMBURGER N, FRITZ H: Effect of human granulocytic elastase on isolated human antithrombin III. *Hoppe-Seyler's Z Physiol Chem* 362:S103-S112, 1981
 111. JOHNSON D, BARRETT AJ, MASON R: Cathepsin L inactivates alpha-1-PI by cleavage of the active site region. *J Biol Chem* 261:14748-14751, 1986
 112. BANDA M, CLARK E, SINHA S, TRAVIS J: Interaction of mouse macrophage elastase with native and oxidized human alpha-1-PI. *J Clin Invest* 79:1314-1317, 1987
 113. DESROCHERS PE, WEISS SJ: Proteolytic inactivation of alpha-1-PI by a neutrophil metalloproteinase. *J Clin Invest* 81:1646-1650, 1988
 114. OKADA Y, WATANABE S, NAKANISHI I, KISHI J, HAYAKAWA T, WATOREK W, TRAVIS J, NAGASE H: Inactivation of tissue inhibitor of metalloproteinases by neutrophil elastase and other serine proteinases. *FEBS J* 229:157-160, 1988
 115. SHIEH B-H, TRAVIS J: The reactive site of human alpha-2-antiplasmin. *J Biol Chem* 262:6055-6059, 1987
 116. OSSANNA PJ, TEST ST, MATHESON NR, REGIANI S, WEISS SJ: Oxidative regulation of neutrophil elastase-alpha-1-PI interactions. *J Clin Invest* 77:1939-1951, 1986
 117. COCHRANE CG, SPRAGG R, REVAG SD: Pathogenesis of adult respiratory distress syndrome: Evidence of oxidant activity in bronchoalveolar lavage fluid. *J Clin Invest* 71:754-761, 1983
 118. MILLER F, KUSCHNER M: Alpha-1-antitrypsin deficiency, emphysema, necrotizing angiitis and glomerulonephritis. *Am J Med* 46:615-623, 1969
 119. MOROZ SP, CUTZ E, BALFE JW, SASS-KORTSAK A: Membranoproliferative glomerulonephritis in childhood cirrhosis associated with alpha-1-antitrypsin deficiency. *Pediatrics* 57:232-238, 1976
 120. RODRIGUEZ-SORIANO J, FIDALGO I, CAMARERO C, VALLO A, OLIVEROS R: Juvenile cirrhosis and membranous glomerulonephritis in a child with alpha-1-antitrypsin deficiency. *Acta Ped Scand* 67:793-796, 1978
 121. LEWIS M, KALLENBACH J, ZALTZMAN M, LEVY H, LURIE D, BAYNES R, KING P, MEYERS A: Severe deficiency of alpha-1-antitrypsin associated with cutaneous vasculitis, rapidly progressive glomerulonephritis, and colitis. *Am J Med* 79:489-494, 1985
 122. SHAH SV: Role of reactive oxygen metabolites in experimental glomerular disease. *Kidney Int* 35:1093-1106, 1989
 123. PEPPIN GJ, WEISS SJ: Activation of endogenous metalloproteinase, gelatinase, by triggered human neutrophils. *Proc Natl Acad Sci USA* 83:4322-4326, 1986
 124. STRICKLIN GP, HOIDAL JR: Oxidant mediated inactivation of TIMP, in *Proceedings of the Matrix Metalloproteinase Conference*, 11-15 September, 1989 Destin, Florida (in press)
 125. CHANG J, WIGLEY F, NEWCOMBE D: Neutral proteinase activation of peritoneal macrophage prostaglandin synthesis. *Proc Natl Acad Sci USA* 77:4736-4740, 1980
 126. ZOJA C, ORISIO S, PERICO N, BENIGNI A, MORIGI M, BENATTI L, RAMBALDI A, REMUZZI G: Constitutive expression of endothelin gene in cultured human mesangial cells and its modulation by transforming growth factor-beta, thrombin, and a thromboxane A₂ analogue. *Lab Invest* 64:16-20, 1991
 127. DOUSA TP, SHAH SV, ABBODD HE: Potential role of cyclic nucleotides in glomerular pathophysiology, in *Advances in Cyclic Nucleotide Research*, edited by HAMET P, SANDS H, New York, Raven Press, 1980, vol 12, pp 285-298
 128. DWORKIN LD, UCHIKAWA I, BRENNER BM: Hormonal modulation of glomerular function. *Am J Physiol* 244:(Renal Fluid Electrol Physiol 13): F95-F104, 1983
 129. IGNARRO LJ: Hormonal control of lysosomal enzyme release from human neutrophils by cyclic nucleotides and autonomic neurohormones, in *Cyclic Nucleotides in Diseases*, edited by WEISS B, Baltimore, University Park, 1975, pp 187-210
 130. LICHTENSTEIN LM: Hormone receptor modulation of cAMP in the control of allergic and inflammatory responses, in *The Role of Immunological Factors in Infections, Allergic, and Antiimmune Processes*, edited by BEERS RF, BASSETT EG, New York, Raven Press, 1976, pp 339-354
 131. ANDERSON WB, JAWORSKI CJ, VLAHAKIS G: Proteolytic activation of adenylate cyclase from cultured fibroblasts. *J Biol Chem* 253:2921-2926, 1978
 132. MOSS J, MANGANIELLO VC, VAUGHAN M: Effects of chymotrypsin and a calcium-dependent protein activator on guanosine 3', 5'-monophosphate phosphodiesterase from rat liver. *Biochim Biophys Acta* 541:279-287, 1978
 133. WALLACH D, ANDERSON W, PASTAN I: Activation of adenylate cyclase in cultured fibroblasts by trypsin. *J Biol Chem* 253:24-26, 1978
 134. RICHERT ND, RYAN RJ: Proteolytic enzyme activation of rat

- ovarian adenylate cyclase. *Proc Natl Acad Sci USA* 74:4857-4861, 1977
135. JOHNSON RA, JAKOBS KH, SCHULTZ G: Extraction of the adenylate cyclase-activating factor of bovine sperm and its identification as a trypsin-like protease. *J Biol Chem* 260:114-121, 1985
 136. SHNEYOUR A, PATT Y, TRAININ N: Trypsin-induced increase in intracellular cyclic AMP of lymphocytes. *J Immunol* 117:2143-2149, 1976
 137. CHEUNG WY: Cyclic 3',5'-nucleotide phosphodiesterase. Evidence for and properties of a protein activator. *J Biol Chem* 246:2859-2869, 1971
 138. SHAH SV: Effect of thrombin on cyclic AMP content in glomeruli isolated from rat kidney. *Kidney Int* 35:824-829, 1989
 139. BABIOR BM: Oxygen-dependent microbial killing by phagocytes. *N Engl J Med* 298:659-668, 1978
 140. KITAGAWA S, TAKAKU F, SAKAMOTA S: Evidence that proteases are involved in superoxide production by human polymorphonuclear leukocytes and monocytes. *J Clin Invest* 65:74-81, 1980
 141. JOHNSON RB JR, CHADWICK DA, COHN ZA: Priming of macrophages for enhanced oxidative metabolism by exposure to proteolytic enzymes. *J Exp Med* 153:1678-1683, 1981
 142. SPEER CP, PABST MJ, HEDEGAARD HB, REST RF, JOHNSTON RB: Enhanced release of oxygen metabolites by monocyte-derived macrophages exposed to proteolytic enzymes: Activity of neutrophil elastase and cathepsin G1. *J Immunol* 133:2151-2156, 1984
 143. SHAH SV: Light emission by isolated rat glomeruli in response to phorbol myristate acetate. *J Lab Clin Med* 98:46-57, 1981
 144. BAUD L, HAGEGE J, SRAER J, RONDEAU E, PEREZ J, ARDAILLOU R: Reactive oxygen production by cultured rat mesangial cells during phagocytosis is associated with stimulation of lipooxygenase activity. *J Exp Med* 158:1836-1852, 1983
 145. JOHNSON RJ, COUSER WG, ALPERS CE, VISSERS M, SCHULZE M, KLEBANOFF SJ: The human neutrophil serine proteinases, elastase and cathepsin G, can mediate glomerular injury in vivo. *J Exp Med* 168:1169-1174, 1988
 146. SCHRIJVER G, SCHALKWIJK J, ROBBEN JCM, ASSMANN KJM, KOENE RAP: Antiglomerular basement membrane nephritis in beige mice. *J Exp Med* 169:1435-1448, 1989
 147. BARICOS WH, SHAH SV: Role of cathepsin B and L in anti-glomerular basement membrane nephritis in rats. *Renal Physiol Biochem* 12:400-405, 1989
 148. BARRETT AJ, KEMBHAVI A, BROWN M, KIRSCHKE H, KNIGHT CG, TAMAI M, HANADA K: L-trans-epoxysuccinyl-leucylamido (4-guanido)butane (E-64) and its analogues as inhibitors of cysteine proteinases including cathepsin B, H and L. *Biochem J* 201:189-198, 1982
 149. TAMAI M, HANADA K, ADACHI T, OGUMA K, KASHIWAGI K, OMURA S, OHZEKI M: Papain inhibitions by optically active E-64 analogs. *J Biochem (Tokyo)* 90:255-257, 1981
 150. HANADA K, TAMAI M, ADACHI T, OGUMA K, KASHIWAGI K, OHMURA S, KOMINAMI E, TOWATARI T, KATUNUMA N: Characterization of the three new analogs of E-64 and their therapeutic application, in *Proteinase Inhibitors: Medical and Biological Aspects*, edited by KATUNUMA N, UMEZAWA H, HOLZER H, Tokyo, Japan Sci. Soc. Press, 1983, p. 25
 151. BARICOS WH, CORTEZ SL, LE QC, WU L-T, SHAW E, HANADA K, SHAH SV: Evidence suggesting a role for cathepsin L in an experimental model of glomerulonephritis. *Arch Biochem Biophys* (in press)
 152. KIRSCHKE H, WIKSTROM P, SHAW E: Active center differences between cathepsins L and B: The S1 binding region. *FEBS Lett* 228:128-130, 1988
 153. MASON RW, GREEN GD, BARRETT AJ: Human liver cathepsin L. *Biochem J* 226:233-241, 1985
 154. SILVER IA, MURRILLS RJ, ETHERINGTON DJ: Microelectrode studies on the acid microenvironment beneath adherent macrophages and osteoclasts. *Exp Cell Res* 175:266-276, 1988
 155. MURACHI T: Intracellular regulatory system involving calpain and calpastatin. *Biochem Int* 18:263-294, 1989
 156. KIRSCHKE H, WIEDRANDERS D, BROMME D, RINNE A: Cathepsin S from bovine spleen. Purification, distribution, intracellular localization and action on proteins. *Biochem J* 264:467-473, 1989
 157. JENNETTE JC, TIDWELL RR, GERATZ JD, BING DH, FALK RJ: Amelioration of immune complex-mediated glomerulonephritis by synthetic protease inhibitors. *Am J Pathol* 127:499-506, 1987
 158. Ikehara S, Shimamura K, Aoyama T, Fujii S, Hamashima Y: Effect of FUT-175, a new synthetic protease inhibitor, on the development of lupus nephritis in (NZB x NZW)F1 mice. *Immunol* 55:595-600, 1985

Appendix. Abbreviations

PMN: polymorphonuclear leukocyte
 GN: glomerulonephritis
 ROM: reactive oxygen metabolites
 APMA: aminophenylmercuric acetate
 SBTI: soya bean trypsin inhibitor
 alpha-1-PI: alpha-1-proteinase inhibitor
 alpha-2-AP: alpha-2-antiplasmin
 PMSF: phenylmethylsulfonylfluoride
 TIMP: tissue inhibitor of metalloproteinases
 tPA: tissue plasminogen activator
 uPA: urokinase-like plasminogen activator
 TLCK: tosyllysyl chloromethyl ketone
 TPCK: tosylphenylalanine chloromethyl ketone
 DIFP: diisopropyl fluorophosphate
 E-64: trans-epoxysuccinyl-L-leucylamido(4-guanidino)butane
 Ep-475: trans-epoxysuccinyl-L-leucylamido-(3-methyl)butane
 SDS: sodium dodecylsulfate
 PMA: phorbol myristate acetate
 PAF: platelet activating factor
 ZFYCHN₂: Z-Phe-Tyr(O-t-butyl)CHN₂
 IP₃: inositol triphosphate